

HORTOPLUS NPK 20.14.15

EC FERTILIZER NPK fertilizer 20.14.15 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

Composition:

Total nitrogen(N).....	20%
Ammoniacal nitrogen.....	4%
nitric nitrogen.....	4%
ureic nitrogen.....	12%
Phosphoric anhydride (P ₂ O ₅) water soluble.....	14%
Potassium oxide(K ₂ O) water soluble.....	15%
Copper (Cu)	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn)	0.05%

Characteristics:

HORTOPLUS 20.14.15 is a water soluble fertilizer made up of pure raw materials.

Its elements ratio makes it ideal in all vegetative phases of the plants.

Its rate of elements makes it particularly suitable in the first vegetative stages and during increasing stage of fruits.

Dose and instructions for use - fertirrigation

Vegetables and strawberry: 5-10 Kgs per 1,000 sm of soil every 8/10 day in the first vegetative stages and during increasing stage of fruits.

Citrus, olive, vine, and actinidia: 70-120 Kgs per hectare during increasing stage of fruits.

Pear, apple, peach, apricot, plum and cherry: 70-120 Kgs per hectare during increasing stage of fruits .

Fodder and industrial crops: 50-100 Kgs per hectare in the first vegetative stages.

Ornamental and flowering plants: 5-10 Kgs per 1,000 sm of soil every 8/10 days in the first vegetative stages.



HORTOPLUS NPK 15.6.30

EC FERTILIZER NPK fertilizer 15.6.30 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

Composition:

Total nitrogen(N).....	15%
Ammoniacal nitrogen.....	5%
Nitric nitrogen.....	5%
Ureic nitrogen.....	5%
Phosphoric anhydride (P ₂ O ₅) water soluble.....	6%
Potassium oxide(K ₂ O) water soluble.....	30%
Copper (Cu)	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn)	0.05%

Characteristics:

HORTOPLUS 15.6.30 is a mineral water soluble fertilizer with high potassium content. It is recommended during increasing stage of fruits and during ripening stage.

The high potassium content determines productions of great value promoting an intense coloration and higher quality of fruits.

Dose and instructions for use - fertirrigation

Vegetables and strawberry: 5-10 Kgs per 1,000 sm of soil every 8/10 days during increasing and ripening stages of the fruits.

Citrus, olive, vine, and actinidia: 70-120 Kgs per hectare during increasing and ripening stages of the fruits.

Pear, apple, peach, apricot, plum and cherry: 70-120 Kgs per hectare during increasing and ripening stages of the fruits.

Fodder and industrial crops: 50-100 Kgs per hectare during vegetative phase.

Ornamental and flowering plants: 5-10 Kgs per 1,000 sm of soil every 8/10 days during vegetative phase.

