

HORTOPLUS NPK 20.20.20

EC FERTILIZER NPK fertilizer 20.20.20 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

Composition:

Total nitrogen(N).....	20%
Ammoniacal nitrogen.....	4%
nitric nitrogen.....	6%
ureic nitrogen.....	10%
Phosphoric anhydride (P ₂ O ₅)water soluble.....	20%
Potassium oxide(K ₂ O) water soluble.....	20%
Copper (Cu)	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn)	0.05%

Characteristics:

HORTOPLUS NPK 20.20.20 is a high water soluble fertilizer.

Its elements ratio makes it ideal in all vegetative phases of the plants.

Microelements presence, besides to intervene how activator for absorption of nitrogen, phosphorus and potassium, has a nutritional role for the plants.

Dose and instructions for use - fertirrigation

Vegetables and strawberry: 5-10 Kgs per 1,000 sm of soil every 8/10 days during all vegetative phases.

Citrus, olive, vine, and actinidia: 70-120 Kgs per hectare during all vegetative phases.

Pear, apple, peach, apricot, plum and cherry: 70-120 Kgs per hectare during all vegetative phases.

Fodder and industrial crops: 50-100 Kgs per hectare during all vegetative phases.

Ornamental and flowering plants: 5-10 Kgs per 1,000 sm of soil every 8/10 days during all vegetative phases.



HORTOPLUS NPK 17.14.23

EC FERTILIZER NPK fertilizer 17.14.23 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

Composition:

Total nitrogen(N).....	17%
Ammoniacal nitrogen.....	5%
Nitric nitrogen.....	6%
Ureic nitrogen.....	6%
Phosphoric anhydride (P ₂ O ₅)water soluble.....	14%
Potassium oxide(K ₂ O) water soluble.....	23%
Copper (Cu)	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn)	0.05%

Charatteristiche:

HORTOPLUS 17.14.23 is a mineral water soluble fertilizer particularly suitable during increasing of the fruits.

Its optimal elements ratio produces a balanced vegetative and productive development.

Dose and instructions for use - fertirrigation

Vegetables and strawberry: 5-10 Kgs per 1,000 sm of soil every 8/10 days during increasing of the fruits.

Citrus, olive, vine, and actinidia: 70-120 Kgs per hectare during increasing of the fruits.

Pear, apple, peach, apricot, plum and cherry: 70-120 Kgs per hectare during increasing of the fruits.

Fodder and industrial crops: 50-100 Kgs per hectare during vegetative phase.

Ornamental and flowering plants: 5-10 Kgs per 1,000 sm of soil every 8/10 days during vegetative phase.

