

## TECNOPLUS NPK 18.9.27

EC FERTILIZER NPK fertilizer 18.9.27 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

### Composition:

Total nitrogen(N).....	18%
Ammoniacal nitrogen.....	2%
nitric nitrogen.....	8%
ureic nitrogen.....	8%
Phosphoric anhydride (P <sub>2</sub> O <sub>5</sub> ) water soluble.....	9%
Potassium oxide(K <sub>2</sub> O) water soluble.....	27%
Copper (Cu) .....	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn) .....	0.05%

### Characteristics:

TECNOPLUS 18.9.27 is a mineral water soluble fertilizer with high potassium content. It is recommended during increasing stage of fruits and during ripening stage.

The high potassium content determines productions of great value promoting an intense coloration and higher quality of fruits.

### Dose and instructions for use - fertirrigation

**Vegetables and strawberry:** 5-10 Kgs per 1,000 sm of soil every 8/10 days during increasing and ripening stages of the fruits.

**Citrus, olive, vine, and actinidia:** 70-120 Kgs per hectare during increasing and ripening stages of the fruits.

**Pear, apple, peach, apricot, plum and cherry:** 70-120 Kgs per hectare during increasing and ripening stages of the fruits.

**Fodder and industrial crops:** 50-100 Kgs per hectare during vegetative phase.

**Ornamental and flowering plants:** 5-10 Kgs per 1,000 sm of soil every 8/10 days during vegetative phase.



Conf. Kg. 25

## TECNOPLUS NPK 8.6.40

EC FERTILIZER NPK fertilizer 8.6.40 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

### Composition:

Total nitrogen(N).....	8%
Ammoniacal nitrogen.....	1%
Nitric nitrogen.....	7%
Phosphoric anhydride (P <sub>2</sub> O <sub>5</sub> ) water soluble.....	6%
Potassium oxide(K <sub>2</sub> O) water soluble.....	40%
Copper (Cu) .....	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn) .....	0.05%

### Characteristics:

TECNOPLUS 8.6.40 is a water soluble fertilizer high potassium content containing microelements. Its composition makes it particularly suitable in the ripening stage of fruits.

High potassium content advances the fruits ripening, at the same time it gives better organoleptic characteristics.

### Dose and instructions for use - fertirrigation

**Vegetables and strawberry:** 5-10 Kgs per 1,000 sm of soil every 8/10 day during ripening stage of fruits.

**Citrus, olive, vine, and actinidia:** 70-120 Kgs per hectare during ripening stage of fruits.

**Pear, apple, peach, apricot, plum and cherry:** 70-120 Kgs per hectare during ripening stage of fruits.

**Fodder and industrial crops:** 50-100 Kgs per hectare in the final stage of vegetative cycle.

**Ornamental and flowering plants:** 5-10 Kgs per 1,000 sm of soil every 8/10 days in the final stage of vegetative cycle.



Conf. Kg. 25