

TECNOPLUS NPK 9.18.27

EC FERTILIZER NPK fertilizer 9.18.27 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

Composition:

Total nitrogen(N).....	9%
Ammoniacal nitrogen.....	3%
nitric nitrogen.....	6%
Phosphoric anhydride (P_2O_5) water soluble.....	18%
Potassium oxide(K_2O) water soluble.....	27%
Copper (Cu)	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn)	0.05%

Characteristics:

TECNOPLUS 9.18.27 is a mineral water soluble fertilizer high purity and solubility particularly suitable during flowering and ripening stages.

Dose and instructions for use - fertirrigation

Vegetables and strawberry: 5-10 Kgs per 1,000 sm of soil every 8/10 days during flowering and ripening stages of the fruits.

Citrus, olive, vine, and actinidia: 70-120 Kgs per hectare during flowering and ripening stages of the fruits.

Pear, apple, peach, apricot, plum and cherry: 70-120 Kgs per hectare during flowering and ripening stages of the fruits.

Fodder and industrial crops: 50-100 Kgs per hectare during final phase of vegetative cycle.

Ornamental and flowering plants: 5-10 Kgs per 1,000 sm of soil every 8/10 days during flowering phase.



Conf. Kg. 25

TECNOPLUS NPK 8.24.24

EC FERTILIZER NPK fertilizer 8.24.24 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

Composition:

Total nitrogen(N).....	8%
Ammoniacal nitrogen.....	5%
Nitric nitrogen.....	3%
Phosphoric anhydride (P_2O_5) water soluble.....	24%
Potassium oxide(K_2O) water soluble.....	24%
Copper (Cu)	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn)	0.05%

Characteristics:

TECNOPLUS 8.24.24 is a water soluble fertilizer with a balanced phospho-potassic activity that favours amino acids and proteins synthesis improving organoleptic characteristics of the fruits. It is particularly suitable in the flowering and ripening stages.

Dose and instructions for use - fertirrigation

Vegetables and strawberry: 5-10 Kgs per 1,000 sm of soil every 8/10 days during flowering and ripening stages of the fruits.

Citrus, olive, vine, and actinidia: 70-120 Kgs per hectare during flowering and ripening stages of the fruits.

Pear, apple, peach, apricot, plum and cherry: 70-120 Kgs per hectare during flowering and ripening stages of the fruits.

Fodder and industrial crops: 50-100 Kgs per hectare during final phase of vegetative cycle.

Ornamental and flowering plants: 5-10 Kgs per 1,000 sm of soil every 8/10 days during flowering phase.



Conf. Kg. 25