

## TECNOPLUS NPK 35.6.9

EC FERTILIZER NPK fertilizer 35.6.9 with copper (Cu), manganese (Mn) and zinc (Zn) to mixing obtained.

### Composition:

Total nitrogen(N).....	35%
Ammoniacal nitrogen.....	1%
Nitric nitrogen.....	2%
ureic nitrogen.....	32%
Phosphoric anhydride (P <sub>2</sub> O <sub>5</sub> ) water soluble.....	6%
Potassium oxide(K <sub>2</sub> O) water soluble.....	9%
Copper (Cu) .....	0.05%
Manganese (Mn).....	0.05%
Zinc (Zn) .....	0.05%

### Characteristics:

TECNOPLUS 35.6.9 is a water soluble fertilizer with high content of nitrogen.

Its ratio between elements makes this product particularly suitable in the first stage of crop cycle and during increasing stage of fruits.

Nitrogen stimulates growing of the plants through compounds formation that have high metabolic significance, how enzymes, amino acids and nucleic acids.

### Dose and instructions for use - fertirrigation

**Vegetables and strawberry:** 5-10 Kgs per 1,000 sm of soil every 8/10 day in the first vegetative stages and during increasing stage of fruits.

**Citrus, olive, vine, and actinidia:** 70-120 Kgs per hectare during increasing stage of fruits.

**Pear, apple, peach, apricot, plum and cherry:** 70-120 Kgs per hectare during increasing stage of fruits.

**Fodder and industrial crops:** 50-100 Kgs per hectare in the first vegetative stages.

**Ornamental and flowering plants:** 5-10 Kgs per 1,000 sm of soil every 8/10 days in the first vegetative stages.



Conf. Kg. 25

## TECNOPLUS NPK 20.5.20

EC FERTILIZER NPK fertilizer 20.5.20 with magnesium (MgO) (2) to mixing obtained.

### Composition:

Total nitrogen(N).....	20%
Ammoniacal nitrogen.....	6%
Ureic nitrogen.....	14%
Phosphoric anhydride (P <sub>2</sub> O <sub>5</sub> ) water soluble.....	5%
Potassium oxide(K <sub>2</sub> O) water soluble.....	20%
Magnesium oxide (MgO) water soluble.....	2%

### Characteristics:

HORTOPLUS 5.5.30 is a water soluble fertilizer high potassium content containing magnesium and microelements. Its composition makes it particularly suitable in the ripening stage of fruits.

High potassium content advances the fruits ripening, at the same time it gives better organoleptic characteristics.

### Dose and instructions for use - fertirrigation

**Vegetables and strawberry:** 5-10 Kgs per 1,000 sm of soil every 8/10 day during ripening stage of fruits.

**Citrus, olive, vine, and actinidia:** 70-120 Kgs per hectare during ripening stage of fruits.

**Pear, apple, peach, apricot, plum and cherry:** 70-120 Kgs per hectare during ripening stage of fruits.

**Fodder and industrial crops:** 50-100 Kgs per hectare in the final stage of vegetative cycle.

**Ornamental and flowering plants:** 5-10 Kgs per 1,000 sm of soil every 8/10 days in the final stage of vegetative cycle.



Conf. Kg. 25